


Strangulation

Visible Signs (may not be present)



Strangulation has only recently been identified as one of the most lethal forms of domestic violence: unconsciousness may occur within seconds and death within minutes. When domestic violence perpetrators choke (strangle) their victims, not only is this a felonious assault, but it may be an attempted homicide. Strangulation is an ultimate form of power and control, where the batterer can demonstrate control over the victim's next breath; having devastating psychological effects or a potentially fatal outcome.

Sober and conscious victims of strangulation will first feel terror and severe pain. If strangulation persists, unconsciousness will follow. Before lapsing into unconsciousness, a strangulation victim will usually resist violently, often producing injuries of their own neck in an effort to claw off the assailant, and frequently also producing injury on the face or hands to their assailant. These defensive injuries may not be present if the victim is physically or chemically restrained before the assault.

Observing Changes

Documentation by photographs sequentially for a period of days after the assault is very helpful in establishing a journal of physical evidence.

Victims should also seek medical attention if they experience difficulty breathing, speaking, swallowing or experience nausea, vomiting, lightheadedness, headache, involuntary urination and/or defecation, especially pregnant victims. A medical evaluation may be crucial in detecting internal injuries and saving a life.

Losing Consciousness

Victims may lose consciousness by any one or all of the following methods: blocking of the carotid arteries in the neck (depriving the brain of oxygen), blocking of the jugular veins (preventing deoxygenated blood from exiting the brain), and closing off the airway, making breathing impossible.



Additional Signs and Symptoms

A larger version of the graphic above which contains detailed signs and symptoms is available for download at <https://www.familyjusticecenter.org/resources/signs-and-symptoms-of-strangulation/>

This project is supported all or in part by
Grant No. 2016-TA-AX-K067 awarded by the
Office on Violence Against Women, U.S. Department of Justice.
The opinions, findings, conclusions, and recommendations
expressed in this publication/program/exhibition are those of
the author(s) and do not necessarily reflect the views of the
Department of Justice, Office on Violence Against Women.



a program of



strangulationtraininginstitute.com

Illustrations and Graphic Design by Yesenia Aceves

Crystal Judson Family Justice Center
718 Court E., Tacoma, WA 98402
Helpline: (253) 798-4166
TDD: (253) 798-4620
www.aplaceofhelp.com

Facts of Strangulation (Choking) That Victims Need to Know

Monitor Your SIGNS

Date & Time

Journal Your Signs

Monitor Your SYMPTOMS

Date & Time

Journal Your Symptoms

Date & Time

Journal Any Other Sensation

Signs of Strangulation

Head- pinpoint red spots (petechiae) on scalp, hair pulled, bump(s), skull fracture, concussion.

Face- red or flushed, petechiae, scratch marks.

Eyes and eyelids- petechiae to the left or right eyeball, bloodshot eyes.

Ear- petechiae (external and/or ear canal), bleeding from ear canal.

Nose- bloody nose, broken nose, petechiae.

Mouth- bruising, swollen tongue, swollen lips, cuts/abrasions.

Under the chin- redness, scratch marks, bruise(s), abrasions.


Neck- redness, scratch marks, fingernail impressions, bruise(s), abrasions, swelling, ligature marks.

Chest and shoulders- redness, scratch marks, bruise(s), abrasions.


Diagrams to Mark Visible Injuries

Use a pen or a marker to indicate any visible signs and/or symptoms.


Front




Under Chin




Right Side



Left Side



Back



Inside Mouth

